

Hyperbolic Coxeter n -polytopes with $n + 2$ facets

P. Tumarkin

Abstract. In this paper, we classify all the hyperbolic non-compact Coxeter polytopes of finite volume combinatorial type of which is either a pyramid over a product of two simplices or a product of two simplices of dimension greater than one. Combined with results of Kaplinskaja [5] and Esselmann [3] this completes the classification of hyperbolic Coxeter n -polytopes of finite volume with $n + 2$ facets.

1 Introduction

Consider a convex polytope P in the n -dimensional hyperbolic space $\mathbb{H}^n$.

A polytope is called a *Coxeter polytope* if its dihedral angles are integer parts of π . Any Coxeter polytope is a fundamental domain of the discrete group generated by the reflections with respect to its facets.

We are interested in Coxeter polytopes of finite volume. In fact, there is no complete classification of such polytopes, but some partial results are obtained. For example, Coxeter polytopes in $\mathbb{H}^3$ are completely characterized by Andreev [1], [2]; all the hyperbolic simplices are classified ([6], [7]).

There are some results about hyperbolic n -polytopes with $n + 2$ facets. Kaplinskaja [5] described all Coxeter simplicial prisms, and Esselmann [3] obtained the classification of compact Coxeter hyperbolic n -polytopes with $n + 2$ facets. In this paper, we complete the classification of Coxeter hyperbolic n -polytopes of finite volume with $n + 2$ facets.

The author is grateful to Prof. R. Kellerhals who brought the work [3] to his attention.

2 Gale diagrams

The essential facts about convex polytopes and Gale diagrams which are used in this paper are quoted below. Proofs and details may be found in [4].

Every combinatorial type of n -polytope with k facets can be represented by a Gale diagram. This consists of m points $a_1, \dots, a_m$ of the unit sphere in $\mathbb{R}^{k-(n+1)}$ centered at the origin and (possibly) the origin a_0 which are labeled according to the following rules:

- 1) Each label is a positive integer, and the sum of labels equals k .
- 2) The points that lie in any open half-space bounded by a hyperplane through the origin have labels whose sum is at least two.

The combinatorial type of a convex polytope can be read off from the Gale diagram in the following way. Each point a_i with label μ_i corresponds to μ_i

facets $f_{i,1}, \dots, f_{i,\mu_i}$ of P . For any subset I of the set of facets of P the intersection of facets $\{f_{j,\gamma} | (j, \gamma) \in I\}$ is a face of P if and only if 0 is contained in the relative interior of $\text{conv}\{a_j | (j, \gamma) \notin I\}$.

By *pyramid* in $\mathbb{H}^n$ we mean a convex hull of a point (apex) and an $(n-1)$ -dimensional polytope which is not a simplex. A polytope P is called a *multipyramid* if P is a pyramid over a pyramid.

Now let P be a n -polytope with $n+2$ facets. In this case the Gale diagram is one-dimensional, and each point in the diagram is equal to -1 , 0 or $+1$. The *multiplicity* of -1 , 0 and $+1$ is defined to be the label of correspondent point in the Gale diagram. Suppose that the multiplicities are p , q and r respectively, $p+q+r = n+2$. There exists a polytope correspondent to a Gale diagram if and only if $p, r \geq 2$ holds.

In the case $q = 0$ the combinatorial type of P is a product of two simplices of dimension $p-1$ and $r-1$. The set of facets is a union of two disjoint sets J_p and J_r of order p and r respectively. Any vertex of P is an intersection of $p-1$ facets from J_p and $r-1$ facets from J_r .

If $q = 1$ then the combinatorial type of P is a pyramid over a product of two simplices. If $q > 1$ then the combinatorial type of P is a multipyramid over a product of two simplices.

Lemma 1. *Let P be a Coxeter polytope in $\mathbb{H}^n$ with $n+2$ facets. Then the combinatorial type of P is either a product of two simplices or a pyramid over a product of two simplices.*

Proof. It is sufficient to prove that any pyramid over a pyramid is not a Coxeter polytope.

Suppose that P is a pyramid over a pyramid P' . Since P' is a pyramid, it is not simple. Hence, the apex A of the pyramid P is an ideal vertex of P . Consider a small horosphere h centered in A . The combinatorial type of the intersection $h \cap P$ coincides with the combinatorial type of P' . At the same time, $h \cap P$ is a compact Euclidean Coxeter $(n-1)$ -polytope. But any compact Euclidean Coxeter polytope is simple. □

Any Coxeter polytope P can be represented by its Coxeter diagram $\Sigma(P)$. Nodes of Coxeter diagram correspond to the facets of P . Two nodes are joined by a $(m-2)$ -fold edge or a m -labeled edge if the corresponding dihedral angle equals $\frac{\pi}{m}$. If the corresponding facets are parallel the nodes are joined by a bold edge, and if they diverge then the nodes are joined by a dotted edge labeled by $\cosh(\rho)$, where ρ is the distance between the facets.

Let Σ be a diagram of order k with nodes $v_1, \dots, v_k$. Define a symmetric $k \times k$ matrix $G(\Sigma)$ with $g_{ii} = 1$, $i = 1, \dots, k$, and $g_{ij} = -\cos(\frac{\pi}{k})$ ($g_{ij} = -1$ or $g_{ij} = -\cosh(\rho)$) if v_i and v_j are joined by k -labeled edge (by a bold edge or by a dotted edge labeled by $\cosh(\rho)$ respectively).

If Σ is a Coxeter diagram of a polytope P then the matrix $G(\Sigma)$ coincides with the Gram matrix of P . See [9] for details.

By the signature and the determinant of a diagram Σ we mean the signature and the determinant of the matrix $G(\Sigma)$.

3 Products of two simplices

In this section, we classify all the non-compact n -polytopes of finite volume with $n + 2$ facets, combinatorial type of which is a product of two simplices of dimension greater than one. For the classification of compact polytopes of this combinatorial type see [3]. All simplicial prisms are listed in [5] (see also [9]).

The result of this section is

Theorem 1. *There exists a unique non-compact Coxeter polytope of finite volume which is combinatorially equivalent to a product of two simplices of dimension greater than one. It is shown in Fig. 1.*

A diagram Σ is called a *Lannér diagram* if any subdiagram of Σ is elliptic, and the diagram Σ is neither elliptic nor parabolic. All Lannér diagrams are classified in [6].

Lemma 2. *Let P be a hyperbolic Coxeter polytope which combinatorial type is a product of two simplices and Σ be a Coxeter diagram of P . Then Σ satisfies the following three conditions:*

- (I) Σ is a union of two disjoint Lannér diagrams L_1 and L_2 .
- (II) For any $v_1 \in L_1$ and $v_2 \in L_2$ a diagram $\Sigma \setminus \{v_1, v_2\}$ is either elliptic or parabolic.
- (III) $\text{Det}(\Sigma) = 0$.

Proof. According to Section 2, Σ is a union of two disjoint diagrams, say L_1 and L_2 , where L_1 consists of facets correspondent to -1 in the Gale diagram of P , and L_2 consists of facets correspondent to 1 . Any vertex of P corresponds to a subdiagram of Σ that is $\Sigma \setminus \{v_1, v_2\}$, where $v_1 \in L_1$, $v_2 \in L_2$. Conversely, for any $v_1 \in L_1$ and $v_2 \in L_2$ a diagram $\Sigma \setminus \{v_1, v_2\}$ corresponds to a vertex of P . Thus, $\Sigma \setminus \{v_1, v_2\}$ is either elliptic or parabolic. Any subdiagram of $\Sigma \setminus \{v_1, v_2\}$ is elliptic. In particular, $L_1 \setminus v_1$ and $L_2 \setminus v_2$ are elliptic for any $v_1 \in L_1$ and $v_2 \in L_2$.

By [9], Th. 3.1, a subdiagram of Σ defines a proper face of P if and only if it is elliptic. Hence, L_1 and L_2 are not elliptic. By [9], Th. 4.1, L_1 and L_2 are not parabolic. Therefore, L_1 and L_2 are Lannér diagrams, and we have proved (I) and (II). Condition (III) follows immediately from the fact that the hyperbolic n -polytope P has more than $n + 1$ facets.

□

Conditions (I) and (II) are satisfied by a big number of Coxeter diagrams. All the diagrams containing parabolic subdiagrams are listed in Table 3. The rest diagrams can not correspond to non-compact polytope. To compute the




Table 1. Diagrams satisfying (I) and (II) and containing parabolic subdiagrams.
White and black vertices compose two disjoint Lannér subdiagrams.

determinants of diagrams shown in the left column we use technical tools from [8] (Prop. 13, Prop. 15 and Table 2). No of these polytopes has $Det = 0$.

A direct computation of other determinants shows that the only Coxeter diagram with $Det = 0$ is one shown in Fig. 1. Denote it by Σ_0 . Denote its nodes as shown in Fig. 1.

Now we shall check the signature of Σ_0 . Since $\Sigma_0 \setminus \{v_1, v_2, v_3\}$ is a Lannér

diagram, the negative inertia index is at least one. $\Sigma_0 \setminus \{v_1, v_4\}$ is elliptic, and the positive inertia index is at least 4. The determinant of Σ_0 equals 0. Thus, the signature is $(4,1,1)$, where the first number is the positive inertia index, the second is the negative inertia index and the third one is the corank of Σ_0 .




Figure 1.

By Theorem 2.1 of [9] for any Coxeter diagram of signature $(n, 1, l)$ there exists a polytope in $\mathbb{H}^n$ correspondent to this diagram. Hence, there exists a Coxeter polytope with the Coxeter diagram Σ_0 . According to [9], Prop. 4.2, this polytope has finite volume.

4 Pyramids

In this section, we classify all the n -polytopes of finite volume with $n + 2$ facets, combinatorial type of which is a pyramid over a product of two simplices.

We need the following

Lemma 3 ([8], Prop. 12). *Let Σ be a Coxeter diagram and v be a node of Σ . Suppose that $\Sigma \setminus v$ is a union of two disjoint parabolic diagrams. Then $\text{Det}(\Sigma) = 0$.*

A diagram Σ is called a *quasi-Lannér diagram* if any subdiagram of Σ is either elliptic or parabolic, and the diagram Σ is neither elliptic nor parabolic. The complete list of quasi-Lannér diagrams can be found in [7].

Lemma 4. *Let P be a hyperbolic Coxeter pyramid over a product of two simplices and Σ be a Coxeter diagram of P . Then Σ satisfies the following three conditions:*

- (I) Σ is a union of two quasi-Lannér diagrams L_1 and L_2 . The intersection of L_1 and L_2 is a unique node v . $L_1 \setminus v$ and $L_2 \setminus v$ are not adjacent.
- (II) The diagrams $L_1 \setminus v$ and $L_2 \setminus v$ are parabolic. Any other subdiagram of L_1 or L_2 is elliptic.
- (III) For any $v_1 \in L_1 \setminus v$ and $v_2 \in L_2 \setminus v$ a diagram $\Sigma \setminus \{v_1, v_2\}$ is either elliptic or parabolic.

Any Coxeter diagram satisfying (I)–(III) determines a hyperbolic Coxeter pyramid over a product of two simplices.

Proof. **1.** Let A be the apex of the pyramid P and v be the node of Σ correspondent to the facet of P opposite to A . Consider a small horosphere h centered in A . The intersection $h \cap P$ is a direct product of two Euclidean simplices. Thus, $\Sigma \setminus v$ is a union of two disjoint parabolic diagrams, say S_1 and S_2 . Clearly, S_1 and S_2 are not adjacent.

Any vertex of P except A corresponds to a subdiagram $\Sigma \setminus \{v_1, v_2\}$, where $v_1 \in S_1$, $v_2 \in S_2$. Hence, $\Sigma \setminus \{v_1, v_2\}$ is either elliptic or parabolic. Since S_1 contains at least two vertices, the diagram $\Sigma \setminus \{S_1, v_2\}$ is a proper subdiagram of $\Sigma \setminus \{v_1, v_2\}$. Thus, it is elliptic.

Denote by L_1 and L_2 subdiagrams $S_1 \cup v$ and $S_2 \cup v$ respectively. We have proved that for any node $u \in L_1$ the diagram $L_1 \setminus u$ is either parabolic (if u coincides with v) or elliptic (otherwise). Hence, L_1 is a quasi-Lannér diagram with the only parabolic subdiagram S_1 . Analogously, L_2 is a quasi-Lannér diagram with the only parabolic subdiagram S_2 .

2. We have shown that conditions (I)–(III) are necessary. Now, suppose that Σ satisfies all the three conditions. The negative inertia index of Σ equals 1, since Σ has a Lannér subdiagram L_1 and the subdiagram $\Sigma \setminus v$ is parabolic. For any $v_1 \in L_1 \setminus v$ and $v_2 \in L_2 \setminus v$ the subdiagram $\Sigma \setminus \{v, v_1, v_2\}$ is elliptic. Hence, the positive inertia index of Σ is at least $n - 1$. By Lemma 3 $\text{Det}(\Sigma) = 0$. Thus, the signature of Σ is either $(n-1, 1, 2)$ or $(n, 1, 1)$. According to [9], Th. 2.1, there exists a Coxeter polytope in $\mathbb{H}^k$ with Coxeter diagram Σ , where k is either $n - 1$ or n .

Suppose that the signature of Σ is $(n-1, 1, 2)$. Then there exists a Coxeter polytope P in $\mathbb{H}^{n-1}$ with Coxeter diagram Σ . This is impossible, since $\Sigma \setminus v$ is a parabolic subdiagram of Σ of rank $n - 1$.

Hence, the signature of Σ equals $(n, 1, 1)$, and Σ determines a Coxeter polytope P in $\mathbb{H}^n$. By Lemmas 1 and 2, Σ determines a hyperbolic Coxeter pyramid over a product of two simplices. This polytope is of finite volume by [9], Prop. 4.2. □

Considering all the quasi-Lannér diagrams with exactly one parabolic subdiagram, we obtain the following

Theorem 2. *All the Coxeter polytopes of finite volume the combinatorial type of which is a pyramid over a product of two simplices are listed in Tables 2–13.*

Remark. Black nodes in Tables 2–13 correspond to the bases of pyramids.




Table 2. Pyramids in $\mathbb{H}^3$.




Table 3. Pyramids in $\mathbb{H}^4$.




Table 4. Pyramids in $\mathbb{H}^5$.





Table 5. Pyramids in $\mathbb{H}^6$.






Table 7. Pyramids in $\mathbb{H}^8$.




Table 8. Pyramids in $\mathbb{H}^9$.




Table 9. Pyramid in $\mathbb{H}^{10}$.




Table 10. Pyramids in $\mathbb{H}^{11}$.




Table 11. Pyramids in $\mathbb{H}^{12}$.




Table 12. Pyramids in $\mathbb{H}^{13}$.




Table 13. Pyramid in $\mathbb{H}^{17}$.

References

- [1] E. M. Andreev, On convex polyhedra in Lobachevskii spaces. Math. USSR Sbornik 10 (1970), 413–440.
- [2] E. M. Andreev, On convex polyhedra of finite volume in Lobachevskii spaces. Math. USSR Sbornik 12 (1970), 255–259.
- [3] F. Esselmann, The classification of compact hyperbolic Coxeter d -polytopes with $d + 2$ facets. Comment. Math. Helvetici 71 (1996), 229–242.
- [4] B. Grünbaum, Convex Polytopes. John Wiley & Sons, 1967.
- [5] I. M. Kaplinskaja, Discrete groups generated by reflections in the faces of simplicial prisms in Lobachevskian spaces. Math. Notes 15 (1974), 88–91.
- [6] F. Lannér, On complexes with transitive groups of automorphisms. Comm. Sem. Math. Univ. Lund 11 (1950), 1–71.
- [7] E. B. Vinberg, Discrete groups generated by reflections in Lobachevskii spaces. Mat. USSR sb. 1 (1967), 429–444.
- [8] E. B. Vinberg, The absence of crystallographic groups of reflections in Lobachevsky spaces of large dimensions. Trans. Moscow Math. Soc. 47 (1985), 75–112.
- [9] E. B. Vinberg, Hyperbolic reflection groups. Russian Math. Surveys 40 (1985), 31–75.